


[Web](#) [Images](#) [Video](#) [News](#) [Maps](#) [more »](#)


[Advanced Scholar Search](#)  
[Scholar Preferences](#)  
[Scholar Help](#)

The "AND" operator is unnecessary -- we include all search terms by default. [\[details\]](#)

[Scholar](#)   [All articles](#) - [Recent articles](#)   [Results 1 - 10 of about 499 for \(DC2 protein OR KIAA0404 pr](#)

### [Protein kinase C involvement in apoptosis](#)

M Lucas, V Sánchez-Margalet - General Pharmacology, 1995 - Elsevier

... dephosphorylation and activation of p34 dc2 does not ... I., Weiss MH and Law RE (1994)

Protein kinase C ... induce apoptosis in human malignant glioma cell lines ...

[Cited by 93](#) - [Related articles](#) - [Web Search](#) - [BL Direct](#) - [All 4 versions](#)

### [High Expression of Doublecortin and KIAA0369 Protein in Fetal Brain Suggests Their Specific Role in ...](#) - [▶amjpathol.org \[HTML\]](#)

M Mizuguchi, J Qin, M Yamada, K Ikeda, S Takashima - American Journal of Pathology, 1999 - ASIP

... cells, and I-23 and Bu17 glioma cells, as ... Protein extraction, sodium dodecyl sulfate polyacrylamide electrophoresis, and ... 1200 for anti-DC1 and anti-DC2, 1:200 ...

[Cited by 39](#) - [Related articles](#) - [Web Search](#) - [BL Direct](#) - [All 4 versions](#)

### [Detection of protein kinase activity specifically activated at metaphase-anaphase transition -](#)

[▶jcb.org \[PDF\]](#)

M Sekimata - The Journal of Cell Biology, 1996 - Rockefeller Univ Press

... and KT34 im- munoreactivities in U251 glioma cells. ... of protein utilizing the principle of protein-dye binding ... mitosis is mediated by p34 cdc2 phosphorylation ...

[Cited by 25](#) - [Related articles](#) - [Web Search](#) - [BL Direct](#) - [All 4 versions](#)

### [Monocyte Chemoattractant Protein-1 \(CCL2\) in Inflammatory Disease and Adaptive Immunity: Therapeutic ...](#)

C Daly, BJ Rollins - Microcirculation, 2003 - Informaworld.com

... Given the fact that CCR2 is a G protein-coupled receptor, it is not ... these mice experience selective abnormalities in the trafficking of DC1 and DC2 cells (ie ...

[Cited by 81](#) - [Related articles](#) - [Web Search](#) - [BL Direct](#) - [All 2 versions](#)

### [... by a Recombinant Listeria monocytogenes Vaccine Targeting Tyrosinase Related Protein-2 and Real-...](#)

RM Prins, KW Bruhn, N Craft, JW Lin, CH Kim, SK ... - Neurosurgery, 2006 - neurosurgery-online.com

... numbers of nucleoprotein and tyrosinase related protein-2-specific ... a tumorigenic subline of the GL26 glioma deficient in ... cells, Dr. Ken Rock for the DC2.4 cells ...

[Cited by 5](#) - [Related articles](#) - [Web Search](#) - [BL Direct](#) - [All 3 versions](#)

### [... genes during permanent focal cerebral ischaemia: characterization of the protein kinase 9b5/MARKL1/...](#)

A Schneider, R Laage, O von Ahsen, A Fischer, M ... - Journal of Neurochemistry, 2004 - pt.wkhealth.com

... org/cgi/data/298/5600/1912/ DC2/1 (Manning ... 9b5/MARK4 kinase, eg in competing with protein interactors. ... of the long variant MARK4l in different glioma cell lines ...

[Cited by 21](#) - [Related articles](#) - [Web Search](#) - [BL Direct](#) - [All 5 versions](#)

### [Growth inhibition of glioma cells transfected with the human beta-interferon gene by liposomes ...](#) - [▶aacrjournals.org \[PDF\]](#)

M Mizuno - Cancer Research, 1990 - AACR

... 4.5 Å±1.5 X IFT5 IU/cell (24.8 Å±3.8 ILJ/mg protein) was found ... E. Block of a glioma cell line in S by interferon ... Science (Washington DC), 2/9: 881-883, 1983 ...

Cited by 61 - Related articles - Web Search - All 3 versions

### Protein Transduction Domains Fused to Virus Receptors Improve Cellular Virus Uptake and Enhance ...

F Kuhnel, B Schulte, T Wirth, N Woller, S Schafers ... - Journal of Virology, 2004 - Am Soc Microbiol

... receptor to serve as molecular adapters between the adenoviral fiber knob protein and negatively ... DC2.4 cells were kindly provided by KL Rock (Worcester, Mass ...

Cited by 9 - Related articles - Web Search - BL Direct - All 6 versions

### Protein structure

SJ Winder, AE Knight, J Kendrick-Jones - Dystrophin: Gene, Protein and Cell Biology, 1997 - books.google.com

... full-length isoforms from skeletal muscle (Koenig et al., 1988) and glioma (Tinsley et al ... the first ~ 250 NH 2-terminal amino acids of each protein, giving an ...

Cited by 10 - Related articles - Web Search

### Immunotherapeutic Targeting of Shared Melanoma-Associated Antigens in a Murine Glioma Model - [aacrjournals.org](http://aacrjournals.org) [HTML]

RM Prins, SK Odesa, LM Liao - Cancer Research, 2003 - AACR

... glioma cells, Dr. Ken Rock for DC2.4 cells ... intracranial; TRP-2, tyrosinase-related protein 2; MAA, melanoma-associated antigen; GAA, glioma-associated antigen ...

Cited by 29 - Related articles - Web Search - BL Direct - All 3 versions

Key authors: M Lucas - C Daly - B Nikolic - V Sánchez-Marg... - E Schneider

Google

Result Page: 1 2 3 4 5 6 7 8 9 10 Next

(DC2 protein OR KIAA0404 protein)

Search

[Google Home](#) - [About Google](#) - [About Google Scholar](#)

©2008 Google